

Noncommutative supergeometry and duality.

Albert Schwarz
University of California at Davis,
Davis, CA 95616 *

February 1, 2008

Abstract

We introduce a notion of Q -algebra that can be considered as a generalization of the notion of Q -manifold (a supermanifold equipped with an odd vector field obeying $\{Q, Q\} = 0$). We develop the theory of connections on modules over Q -algebras and prove a general duality theorem for gauge theories on such modules. This theorem contains as a simplest case $SO(d, d, \mathbf{Z})$ -duality of gauge theories on noncommutative tori.

It was shown recently that noncommutative geometry is quite useful in the study of string theory/M-theory (see [4]-[10] and references therein). It was proved, in particular, that the gauge theory on noncommutative tori has $SO(d, d, \mathbf{Z})$ duality group, closely related to T-duality in string theory [5]. The goal of present paper is to derive a very general duality theorem, containing $SO(d, d, \mathbf{Z})$ -duality as a special case. This theorem is formulated and proved in the framework of "noncommutative supergeometry". The main idea of noncommutative geometry is to consider every associative algebra as an algebra of functions on "noncommutative space". Of course, supergeometry fits very nicely in this approach: one of the most convenient definitions of a supermanifold is formulated in terms of the algebra of functions on it. One can say that supergeometry is "supercommutative $\mathbf{Z}_2$ -graded noncommutative geometry".

One of important notions of supergeometry is the notion of Q -manifold (of a manifold equipped with an odd vector field Q satisfying $\{Q, Q\} = 0$); see [12]. The first order differential operator $\hat{Q}$

*Partially supported by NSF grant DMS-9801009

corresponding to Q obeys $\hat{Q}^2 = 0$; therefore the algebra of functions on a Q -manifold can be considered as a differential $\mathbf{Z}_2$ -graded associative algebra and it is naturally to think that differential $\mathbf{Z}_2$ -graded associative algebras are analogs of Q -manifolds. However, in this paper we introduce another notion, the notion of Q -algebra, that also can be considered as a natural generalization of Q -manifold and that can be used to develop the theory of connections and to prove a general duality theorem. Namely, we define a Q -algebra as a $\mathbf{Z}_2$ -graded associative algebra Ω equipped with an odd derivation Q obeying $Q^2 a = [\omega, a]$; here $\omega \in \Omega$ should satisfy $Q\omega = 0$. (One says that a linear operator acting on graded algebra is a derivation, if it satisfies the graded Leibniz rule.) Of course, in the case when Ω is supercommutative this definition coincides with the definition of differential algebra, but if we do not assume supercommutativity this definition is more general. We define a connection on Ω -module $\mathcal{E}$ as a map $\nabla : \mathcal{E} \rightarrow \mathcal{E}$ obeying the Leibniz rule $\nabla(ea) = \nabla e \cdot a + (-1)^{\deg e} e \cdot Qa$; the general duality theorem is formulated in terms of such connections. We analyze the relation of the standard definition of connection in noncommutative geometry to our one. It seems that many well known constructions and theorems become more transparent in our formalism. From the other side many considerations of present paper are similar to arguments employed previously, especially in [1], [2], [3].

Preliminaries.

When we talk about associative algebra Ω we always have in mind graded ($\mathbf{Z}$ -graded or $\mathbf{Z}_2$ -graded) unital associative algebra over $\mathbf{C}$. Graded commutator is defined by the formula

$$[a, b] = ab - (-1)^{\deg a \cdot \deg b} ba.$$

In what follows all commutators are understood as graded commutators.

A (right) module $\mathcal{E}$ over Ω is a graded vector space with operator of multiplication on elements of Ω from the right; this operation should have standard properties: $(ea) \cdot b = e \cdot (ab)$ $e(a+b) = ea + eb$ etc. Grading on $\mathcal{E}$ should be compatible with grading on Ω (i.e. $\deg(ea) = \deg e + \deg a$). The definition of a left module is similar; by default our modules are right modules.

A vector space $\mathcal{E}$ is called an (Ω_1, Ω_2) -bimodule if it is a left Ω_1 -module and a right Ω_2 -module; we require that $(a_1 e) a_2 = a_1 (e a_2)$

where $a_i \in \Omega_i$, $e \in \mathcal{E}$.

If $\mathcal{E}_1, \mathcal{E}_2$ are Ω -modules we define an Ω -homomorphism as a map $\varphi : \mathcal{E}_1 \rightarrow \mathcal{E}_2$ obeying $\varphi(xa) = \varphi(x)a$. The graded algebra of Ω -homomorphisms of the Ω -module $\mathcal{E}$ into itself (algebra of Ω -endomorphisms) is denoted by $\text{End}_\Omega \mathcal{E}$. If $\mathcal{E}$ is an (Ω_1, Ω_2) -bimodule there exist natural homomorphisms $\Omega_1 \rightarrow \text{End}_{\Omega_2} \mathcal{E}$ and $\Omega_2 \rightarrow \text{End}_{\Omega_1} \mathcal{E}$.

If $\mathcal{E}_1$ is a right Ω -module and $\mathcal{E}_2$ is a left Ω -module we define $\mathcal{E}_1 \otimes_\Omega \mathcal{E}_2$ as a vector space obtained from the standard tensor product $\mathcal{E}_1 \otimes_{\mathbf{C}} \mathcal{E}_2$ by means of identification $e_1 a \otimes e_2 \sim e_1 \otimes a e_2$, where $e_i \in \mathcal{E}_i$, $a \in \Omega$.

We say that (Ω_1, Ω_2) -bimodule P_1 and (Ω_2, Ω_1) -bimodule P_2 are Morita equivalence bimodules if $P_1 \otimes_{\Omega_2} P_2$ is isomorphic to Ω_1 as (Ω_1, Ω_1) -bimodule and $P_2 \otimes_{\Omega_1} P_1$ is isomorphic to Ω_2 as (Ω_2, Ω_2) -bimodule. If such bimodules P_1, P_2 exist we say that Ω_1 and Ω_2 are Morita equivalent. For every Ω_1 -module we can define an Ω_2 -module $\tilde{\mathcal{E}} = \mathcal{E} \otimes_{\Omega_1} P_1$. Applying an analogous construction to Ω_2 -module $\tilde{\mathcal{E}}$ and bimodule P_2 we obtain an inverse map: $\mathcal{E} = \tilde{\mathcal{E}} \otimes_{\Omega_2} P_2$. One can say that Morita equivalent algebras have equivalent categories of modules.

A (finitely generated) free module Ω^n over Ω can be defined as the space of column vector with entries from Ω and with componentwise multiplication on elements of Ω . We regard Ω^n as a right module, but it can be considered also as (Ω, Ω) -bimodule. (We already used the structure of (Ω, Ω) -bimodule on $\Omega^1 = \Omega$.) The algebra $\text{End}_\Omega \Omega^n$ of endomorphisms of Ω^n can be identified the algebra $\text{Mat}_n \Omega$ of $n \times n$ matrices with entries from Ω ; these matrices act on Ω^n by means of multiplication from the left. A projective Ω -module can be defined as a direct summand E in a free module Ω^n . The decomposition $\Omega^n = E + E'$ into a direct sum determines an endomorphism $e : \Omega^n \rightarrow \Omega^n$ projecting Ω^n onto E ; in other words $e^2 = e$, $ex = x$ for $x \in E$, $ex' = 0$ for $x' \in E'$. Notice that in our terminology projective modules are always finitely generated.

Projective Ω -modules form a semigroup with respect to direct summation. Applying the Grothendieck construction to this semigroup we obtain the K-theory group $K_0(\Omega)$. More precisely, we say that a projective module E specifies an element $[E] \in K_0(\Omega)$ and impose the relation $[E_1 + E_2] = [E_1] + [E_2]$. A $\mathbf{C}$ -linear map $\tau : \Omega \rightarrow \mathbf{C}$ is called a (graded) trace if it vanishes on all (graded) commutators: $\tau([a, b]) = 0$ for all $a, b \in \Omega$. We always consider

graded traces; therefore we almost always omit the word "graded" in our formulations.

A trace τ on Ω generates a trace on $\text{End}_\Omega \Omega^n = \text{Mat}_n \Omega$; this trace will be denoted by the same symbol τ . (To calculate the trace of a matrix $(a_{ij}) \in \text{Mat}_n \Omega$ one should take the supertrace of the matrix $(\tau(a_{ij}))$).

If $E \subset \Omega^n$ is a projective module then the algebra $\text{End}_\Omega E$ of endomorphisms of E can be identified with the subalgebra of $\text{End}_\Omega \Omega^n = \text{Mat}_n \Omega$ consisting of elements of the form eae . (Here $e : \Omega^n \rightarrow \Omega^n$ is a projection of Ω^n onto E , $a \in \text{End}_\Omega \Omega^n$). We define a graded trace $\tilde{\tau}$ on $\text{End}_\Omega E$ as a restriction of τ to this subalgebra.

If $\mathcal{E}$ is an Ω -module, then starting with an element $g \in \Omega$ and Ω -homomorphism $f : \mathcal{E} \rightarrow \Omega$ we can construct an endomorphism $g \otimes f : \mathcal{E} \rightarrow \mathcal{E}$ transforming $x \in \mathcal{E}$ into $gf(x) \in \mathcal{E}$. (The endomorphism $g \otimes f$ can be considered as a generalization of linear operator of rank 1.)

For any algebra Ω we construct a vector space $\bar{\Omega} = \Omega/[\Omega, \Omega]$ factorizing the vector space Ω with respect to the subspace $[\Omega, \Omega]$ spanned by all (graded) commutators $[a, b]$. This construction is closely related with the notion of trace: traces on Ω correspond to linear functionals on $\bar{\Omega}$.

If $\mathcal{E}$ is a projective Ω -module, one can construct a $\mathbf{C}$ -linear map $\text{Tr} : \text{End}_\Omega \mathcal{E} \rightarrow \bar{\Omega}$ transforming an endomorphism of the form $g \otimes f$ into the class $\overline{f(g)} \in \bar{\Omega}$ of $f(g) \in \Omega$. (Such a $\mathbf{C}$ -linear map is unique because in the case of projective module every endomorphism can be represented as a finite sum of endomorphisms of the form $g \otimes f$.) The map Tr has the main property of trace

$$\text{Tr}[\varphi, \psi] = 0$$

(trace of graded commutator of two Ω -endomorphisms $\varphi, \psi \in \text{End}_\Omega \mathcal{E}$ vanishes). In some sense the map $\text{Tr} : \text{End}_\Omega \mathcal{E} \rightarrow \bar{\Omega}$ can be considered as universal trace on $\text{End}_\Omega \mathcal{E}$. (As we mentioned every trace τ on Ω determines a trace $\tilde{\tau}$ on $\text{End}_\Omega \mathcal{E}$. It is easy to verify that $\tilde{\tau}(\varphi) = \tau(\text{Tr}\varphi)$.)

Q -algebras

Definition. Let Ω be a graded associative algebra. We say the Ω is a Q -algebra if it is equipped with derivation Q of degree 1 and there exists an element $\omega \in \Omega^2$ satisfying

$$Q^2 x = [\omega, x] \tag{1}$$

for all $x \in \Omega$.

Calculating Q^3x in two ways we obtain

$$Q^3x = Q([\omega, x]) = Q\omega \cdot x + \omega \cdot Qx - Qx \cdot \omega - x \cdot Q\omega \cdot (-1)^{\deg x} \quad (2)$$

$$Q^3x = Q^2 \cdot Qx = [\omega, Qx] = \omega \cdot Qx - Qx \cdot \omega. \quad (3)$$

We see that $Q\omega \cdot x = x \cdot Q\omega \cdot (-1)^{\deg x}$, i.e.

$$[Q\omega, x] = 0$$

We proved that $Q\omega \in \Omega^3$ commutes with all elements of Ω (in the sense of superalgebra). In almost all interesting cases it follows from this condition that $Q\omega$ vanishes.

We will include the additional condition

$$Q\omega = 0$$

in the definition of Q -algebra.

We always consider unital algebras. It is easy to check that applying Q to the unit we get 0. (This follows from the Leibniz rule.)

Let us consider a (graded) Ω -module E . We define a connection on $\mathcal{E}$ as a $\mathbf{C}$ -linear operator $\nabla : E \rightarrow \mathcal{E}$ having degree 1 and obeying the Leibniz rule:

$$\nabla(xa) = (\nabla x) \cdot a + (-1)^{\deg x} \cdot x \cdot Qa. \quad (4)$$

for all $x \in E$, $a \in \Omega$.

Let us introduce the notation

$$\hat{a}x = (-1)^{\deg x \cdot \deg a} xa$$

The formula (4) can be rewritten in the form

$$[\nabla, \hat{a}] = \widehat{Qa}$$

It is easy to check that some standard statements about connections remain true in our case. However, the definition of curvature should be modified.

1) If ∇ is a fixed connection on $\mathcal{E}$, then every other connection has the form

$$\nabla' = \nabla + A$$

where $A \in \text{End}_\Omega^1 \mathcal{E}$ is an arbitrary endomorphism of degree 1.

2) If $\varphi \in \text{End}_\Omega \mathcal{E}$ is an endomorphism then $[\nabla, \varphi]$ is also endomorphism.

3) The operator $\nabla^2 + \hat{\omega}$ is an endomorphism: $\nabla^2 + \hat{\omega} \in \text{End}_\Omega^2 \mathcal{E}$. This endomorphism is called the curvature of connection ∇ ; it is denoted by $F(\nabla)$ (or simply by F). It obeys $[\nabla, F] = 0$.

To check this statement we represent ∇^2 as $\frac{1}{2}[\nabla, \nabla]$ and calculate $[[\nabla, \nabla], \hat{a}]$ by means of (4) and Jacobi identify.

4) Let us define the operator $\tilde{Q} : \text{End}_\Omega \mathcal{E} \rightarrow \text{End}_\Omega \mathcal{E}$ by the formula

$$\tilde{Q}\varphi = [\nabla, \varphi].$$

It is easy to verify that

$$\tilde{Q}^2\varphi = [F, \varphi], \quad (5)$$

where F is the curvature of ∇ .

It follows from this statement and from $\tilde{Q}F = [\nabla, F] = 0$ that the algebra $\text{End}_\Omega \mathcal{E}$ equipped with the operator $\tilde{Q}$ is a Q -algebra with $\tilde{\omega} = F$. (One should notice, however, that we can also take $\tilde{\omega} = F + c$, where c is a central element obeying $\nabla c = 0$.)

Let us consider an (Ω_1, Ω_2) -bimodule P where Ω_1 is a Q -algebra with respect to the operator Q_1 and Ω_2 is a Q -algebra with respect to the operator Q_2 . We say that an operator $\nabla_P : P \rightarrow P$ is a connection on bimodule P if

$$\nabla_P(ax) = (-1)^{\deg a} \cdot a \nabla_P(x) + Q_1 a \cdot x$$

$$\nabla_P(xb) = \nabla_P x \cdot b + (-1)^{\deg x} \cdot x Q_2 b$$

for all $x \in P$, $a \in \Omega_1$, $b \in \Omega_2$.

In other words, ∇_P should be a connection with respect to Ω_1 and with respect to Ω_2 at the same time.

It follows from the above statements that every Ω -module $\mathcal{E}$ equipped with a connection ∇ can be considered as $(\text{End}_\Omega \mathcal{E}, \Omega)$ -bimodule and ∇ is a connection on this bimodule.

Using an (Ω_1, Ω_2) -bimodule P we can assign to every (right) Ω_1 -module $\mathcal{E}$ a (right) Ω_2 -module $\tilde{\mathcal{E}}$ taking the tensor product with P :

$$\tilde{\mathcal{E}} = \mathcal{E} \otimes_{\Omega_1} P \quad (6)$$

(To take the tensor product over Ω_1 we identify $ea \otimes p$ with $e \otimes ap$ in the standard tensor product $\mathcal{E} \otimes_{\mathbf{C}} P$. Here $e \in \mathcal{E}$, $p \in P$, $a \in \Omega_1$.)

If we have a connection ∇_P in the bimodule P we can transfer a connection on $\mathcal{E}$ to a connection on $\tilde{\mathcal{E}}$. Namely, for every connection ∇ on $\mathcal{E}$ we define an operator $\nabla \otimes 1 + 1 \otimes \nabla_P$ on $\mathcal{E} \otimes_{\mathbf{C}} P$. It is easy to check that this operator is compatible with identification $ea \otimes p \sim e \otimes ap$ and therefore descends to an operator $\tilde{\nabla} : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{E}}$. The operator $\tilde{\nabla}$ can be considered as a connection on Ω_2 -module $\tilde{\mathcal{E}}$.

It is easy to relate the curvatures of the connections ∇ and $\tilde{\nabla}$. We should take into account that correspondence between $\mathcal{E}$ and $\tilde{\mathcal{E}}$ is natural, i.e. to every endomorphism $\sigma \in \text{End}_{\Omega_1} \mathcal{E}$ we can assign an endomorphism $\tilde{\sigma} \in \text{End}_{\Omega_2} \tilde{\mathcal{E}}$ (the map $\sigma \otimes 1 : \mathcal{E} \otimes_{\mathbf{C}} P \rightarrow \mathcal{E} \otimes_{\mathbf{C}} P$ descends to an endomorphism $\tilde{\sigma} : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{E}}$). In particular, the curvature $F(\nabla) \in \text{End}_{\Omega_1} \mathcal{E}$ determines an endomorphism $\widetilde{F(\nabla)} \in \text{End}_{\Omega_2} \tilde{\mathcal{E}}$. One can verify that the curvature $F(\tilde{\nabla})$ of the connection $\tilde{\nabla}$ on $\tilde{\mathcal{E}}$ can be represented in the form:

$$F(\tilde{\nabla}) = \widetilde{F(\nabla)} + \tilde{\varphi}, \quad (7)$$

where $\tilde{\varphi}$ is a fixed element of $\text{End}_{\Omega_2} \tilde{\mathcal{E}}$.

To verify (7) we notice that

$$\nabla^2 \otimes 1 + \hat{\omega}_1 \otimes 1 : \mathcal{E} \otimes_{\mathbf{C}} P \rightarrow \mathcal{E} \otimes_{\mathbf{C}} P$$

descends to the endomorphism $\widetilde{F(\nabla)} : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{E}}$ and $\nabla^2 \otimes 1 + 1 \otimes \nabla_P^2 + 1 \otimes \hat{\omega}_2$ descends to $F(\tilde{\nabla}) : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{E}}$. Using the relation $\hat{\omega}_1 \otimes 1 = -1 \otimes \hat{\omega}_1$ we obtain that the map $\tilde{\varphi} = F(\tilde{\nabla}) - \widetilde{F(\nabla)}$ is induced by the map $\varphi = 1 \otimes \psi : \mathcal{E} \otimes_{\mathbf{C}} P \rightarrow \mathcal{E} \otimes_{\mathbf{C}} P$ where the map $\psi : P \rightarrow P$ is given by the formula

$$\psi = \nabla_P^2 + \hat{\omega}_1 + \hat{\omega}_2.$$

It is easy to check that

$$\psi \in \text{End}_{\Omega_1} P \cap \text{End}_{\Omega_2} P \quad (8)$$

(i.e. $\psi(ax) = a\psi(x)$, $\psi(xb) = \psi(x)b$ for $x \in P$, $a \in \Omega_1$, $b \in \Omega_2$). To check that ψ commutes with $a \in \Omega_1$ we represent it in the form $\psi = \widetilde{F_1(\nabla_P)} + \hat{\omega}_2$, where $F_1(\nabla_P)$ stands for the curvature of ∇_P considered as Ω_1 -connection; the representation $\psi = \widetilde{F_2(\nabla_P)} + \hat{\omega}_1$ should be used to prove that $\psi \in \text{End}_{\Omega_2} P$.

It follows from (8) that $\varphi = 1 \otimes \psi$ descends to $\tilde{\mathcal{E}}$ and gives an Ω_2 -endomorphism $\tilde{\varphi}$. One should notice that these facts are clear also from the representation $\tilde{\varphi} = F(\tilde{\nabla}) - \widetilde{F(\nabla)}$.

To illustrate the above statements we can start with an arbitrary Q -algebra Ω and arbitrary Ω -module P with connection ∇_P . We consider P as (Ω_1, Ω_2) -bimodule, where $\Omega_1 = \text{End}_\Omega P$, $\Omega_2 = \Omega$. (We have seen that $\Omega_1 = \text{End}_\Omega P$ is a Q -algebra with respect to the operator $\tilde{Q}\varphi = [\nabla_P, \varphi]$ and that ∇_P is a connection also with respect to this Q -algebra.) It follows from our calculations that $F = F_2(\nabla_P) = \nabla_P^2 + \hat{\omega}_2$, $\hat{\omega}_1 = -F$ and therefore $\psi = F + \hat{\omega}_1 = 0$. (We can obtain the same result noticing that $F_1(\nabla_2) = \nabla_P^2 + \hat{\omega}_1 = (F - \hat{\omega}_1 = -\hat{\omega}_2)$.) We see that in our situation $\varphi = 0$; hence, $F(\tilde{\nabla}) = \widetilde{F(\nabla)}$. (However, as we noticed above one can modify the definition of Q -algebra $\text{End}_\Omega P$ adding central element c with $\nabla c = 0$ to ω_1 ; then $\varphi \neq 0$.)

We would like to give conditions when gauge theories on Ω_1 -module $\mathcal{E}$ and in Ω_2 -module $\tilde{\mathcal{E}}$ are equivalent. To establish such an equivalence we need (Ω_1, Ω_2) -bimodule P' equipped with connection $\nabla_{P'}$. Such a bimodule permits us to transfer modules and connections in opposite direction. If the constructions obtained by means of P' are inverse to constructions specified by P we say that bimodules P, P' give Morita equivalence of Q -algebras Ω_1 and Ω_2 (or that they are Morita equivalence bimodules). Of course, this notion generalizes the standard notion of Morita equivalence of associative algebras, when we do not use the operator Q and connections. The definition of Morita equivalence bimodules can be reformulated in the following more constructive way. Let us suppose that there exist two bilinear scalar products between P and P' taking values in Ω_1 and in Ω_2 respectively. We assume that scalar products are Ω_2 -invariant and Ω_1 -invariant correspondingly. In other words, we assume that for $p \in P$, $p' \in P'$ we have $\langle p, p' \rangle_1 \in \Omega_1$, $\langle p', p \rangle_2 \in \Omega_2$ and $\langle p\omega, p' \rangle_1 = \langle p, \omega p' \rangle_1$ for $\omega \in \Omega_2$, $\langle p' \sigma_1, p \rangle_2 = \langle p', \sigma_1 p \rangle_2$ for $\sigma_1 \in \Omega_1$. We require also that

$$\sigma_1 \langle p, p' \rangle_1 \sigma_2 = \langle \sigma_1 p, p' \sigma_2 \rangle_1, \quad \omega_1 \langle p', p \rangle_2 \omega_2 = \langle \omega_1 p', p \omega_2 \rangle_2 \quad (9)$$

$$p_1 \langle p, p' \rangle_1 = \langle p_1, p \rangle_2 p', \quad \langle p', p \rangle_2 p'_1 = p' \langle p, p'_1 \rangle_1 \quad (10)$$

Here $p, p_1 \in P$, $p', p'_1 \in P'$, $\sigma_i \in \Omega_1$, $\omega_i \in \Omega_2$. The scalar products determine maps

$$\alpha : P \otimes_{\Omega_2} P' \rightarrow \Omega_1, \quad \beta : P' \otimes_{\Omega_1} P \rightarrow \Omega_2.$$

We can consider $P \otimes_{\Omega_2} P'$ and Ω_1 as (Ω_1, Ω_1) -bimodules; then it

follows from (9), that α is a homomorphism of bimodules; similarly β is a homomorphism of (Ω_2, Ω_2) -bimodules. We require that α and β be isomorphisms. Then

$$(\mathcal{E} \otimes_{\Omega_1} P) \otimes_{\Omega_2} P' = \mathcal{E} \otimes_{\Omega_1} (P \otimes_{\Omega_2} P') = \mathcal{E} \otimes_{\Omega_1} \Omega_1 = \mathcal{E}$$

for every Ω_1 -module $\mathcal{E}$. This statement together with similar statement for Ω_2 -modules gives us one-to-one correspondence between Ω_1 -modules and Ω_2 -modules (more precisely it gives us equivalence of categories of Ω_1 -modules and Ω_2 -modules). To obtain one-to-one correspondence between connections we should impose additional requirements

$$\begin{aligned} < \nabla_P p, p' >_1 + < p, \nabla_{P'} p' >_1 = Q < p, p' >_1, \\ < \nabla_{P'} p', p >_2 + < p', \nabla_P p >_2 = Q < p', p >_2 \end{aligned} \quad (11)$$

It follows from our assumptions that the operator

$$\nabla_P \otimes 1 + 1 \otimes \nabla_{P'}$$

on $P \otimes_{\mathbb{C}} P'$ descends to operator Q on $P \otimes_{\Omega_2} P'$. Using that $Q \cdot 1 = 0$ we obtain that the operator $\nabla \otimes 1 + 1 \otimes Q$ on $\mathcal{E} \otimes_{\mathbb{C}} \Omega_1$ descends to ∇ on $\mathcal{E} \otimes_{\Omega_1} \Omega_1 = \mathcal{E}$. This means that going from Ω_1 -connections to Ω_2 -connections and back we obtain the original Ω_1 -connection. This fact together with similar statement about Ω_2 -connections gives one-to-one correspondence between Ω_1 -connections and Ω_2 -connections.

We see that under our conditions we have equivalence between gauge theories on Ω_1 -module $\mathcal{E}$ and on Ω_2 -module $\tilde{\mathcal{E}}$ (duality). We will describe later how the duality of gauge theories on noncommutative tori can be obtained this way.

Let us study connections on projective Ω -modules where Ω is a Q -algebra.

First of all it is easy to construct a connection on an arbitrary projective Ω -module $\mathcal{E}$. Namely, if $\mathcal{E}$ is specified by means of projection $e : \Omega^n \rightarrow \Omega^n$ (i.e. $e\Omega^n = \mathcal{E}$) we can construct a connection on $\mathcal{E}$ (so called Levi-Civita connection) by means of the formula $D = eQe$ where Q acts on Ω^n componentwise. (The Leibniz rule for D follows from $e^2 = e$ and from the Leibniz rule for Q .) The curvature of the Levi-Civita connection is given by the formula:

$$F = e((Qe)^2 + \omega \cdot 1).$$

For any algebra Ω we defined a vector space $\bar{\Omega} = \Omega/[\Omega, \Omega]$. If Ω is a Q -algebra we have $Q([\Omega, \Omega]) \subset [\Omega, \Omega]$. This means that the operator $Q : \Omega \rightarrow \Omega$ descends to an operator $\bar{Q} : \bar{\Omega} \rightarrow \bar{\Omega}$. It is easy to check that $\bar{Q}$ is a differential: $\bar{Q}^2 = 0$.

Now we will define the Chern character of a connection D on a projective Ω -module $\mathcal{E}$ as an element of $\bar{\Omega}$:

$$\text{ch}D = \sum_{q=0} \frac{1}{q!} \text{Tr} F^q$$

(Recall that we defined a map $\text{Tr} : \text{End}_{\Omega}\mathcal{E} \rightarrow \bar{\Omega}$ using the formula $\text{Tr}(g \otimes f) = \overline{f(g)}$. Here $f : \mathcal{E} \rightarrow \Omega$ is an Ω -homomorphism, $g \in \Omega$ and $g \otimes f$ transforms $x \in \mathcal{E}$ into $gf(x) \in \mathcal{E}$. The map $a \rightarrow \bar{a}$ transforms $a \in \Omega$ into its class $\bar{a} \in \bar{\Omega}$.)

One can prove the following statements:

1) $\text{ch}D$ is closed with respect to the differential $\bar{Q}$ in $\bar{\Omega}$:

$$\bar{Q}\text{ch}D = 0 \tag{12}$$

2) If D', D are two connections on Ω -module $\mathcal{E}$ then $\text{ch}D' - \text{ch}D$ is exact with respect to the differential $\bar{Q}$:

$$\text{ch}D' - \text{ch}D = \bar{Q}(\text{something}). \tag{13}$$

The proof is based on the following lemma:

For every endomorphism $\varphi \in \text{End}_{\Omega}\mathcal{E}$ we have

$$\text{Tr}[D, \varphi] = \bar{Q}\text{Tr}\varphi \tag{14}$$

It is sufficient to verify (14) for Levi-Civita connection $D = eQe$ (because $\text{Tr}[D' - D, \varphi] = 0$) and for endomorphisms of the form $\varphi = g \otimes f$ (because these endomorphisms span $\text{End}_{\Omega}\mathcal{E}$).

Using (14) we deduce (12) from the relation $[D, F^q] = 0$ that follows immediately from $[D, F] = 0$.

To derive (13) we will consider a smooth family $D(t) = D + t(D' - D)$ of connections on $\mathcal{E}$ and prove that

$$\frac{d}{dt}\text{ch}D(t) = \bar{Q}(\text{something}).$$

First of all we notice that the curvature $F(t)$ of connection $D(t)$ obeys

$$\frac{dF(t)}{dt} = [\Gamma, D(t)]$$

where $\Gamma = D' - D \in \text{End}_\Omega \mathcal{E}$. We see that

$$\frac{dF}{dt} = [\Gamma, D] \mod [\bar{\Omega}, \bar{\Omega}].$$

and therefore

$$\frac{dF^q}{dt} = q[\Gamma, D]F^{q-1} = q[D, \Gamma F^{q-1}] \mod [\bar{\Omega}, \bar{\Omega}],$$

$$\frac{d\text{Tr} F^q}{dt} = q\text{Tr}[D, \Gamma F^{q-1}] \in \bar{Q}(\bar{\Omega}).$$

Integrating over t we obtain (13).

In the proof of (13) we assumed that Ω is equipped with topology having some properties that permit us to justify the calculations above. These assumptions are not necessary; it is easy to modify our consideration to obtain completely algebraic proof (as in [2] for example).

Sometimes it is convenient to reformulate (13) using the notion of closed trace. We say that a linear functional on Ω is a closed trace if it vanishes on (graded) commutators and on elements of the form Qa . It follows from (13) that for a closed trace τ the number $\tau(\text{ch}(D))$ does not depend on the choice of the connection D on the module $\mathcal{E}$; it depends only on the K -theory class of the module $\mathcal{E}$.

Using the differential $\bar{Q}$ we can define the homology $H(\bar{\Omega})$ in the standard way: $H(\bar{\Omega}) = \ker \bar{Q} / \text{Im} \bar{Q}$. It follows from (12), (13) that the Chern character specifies a homomorphism $\text{ch}: K_0(\Omega) \rightarrow H^{\text{even}}(\bar{\Omega})$. One can construct also a map $K_1(\Omega) \rightarrow H^{\text{odd}}(\bar{\Omega})$; we will not discuss this construction here.

Connections on modules over associative algebras.

The theory of connections on modules over Q -algebras can be considered as a generalization of the theory of connections on associative algebras. If A is an associative algebra one can construct a differential $\mathbf{Z}$ -graded algebra $\Omega(A) = \sum_{n \geq 0} \Omega^n(A)$ (universal differential graded algebra) in the following way. The vector space $\Omega^n(A)$ is spanned by formal expressions $a_0 da_1 \dots da_n$ and $\lambda da_1 \dots da_m$ where $a_0, \dots, a_n \in A$, $n \geq 0$, $m \geq 1$, $\lambda \in \mathbf{C}$. The multiplication and the differential on $\Omega(A)$ are defined by means of Leibniz rule and relation $d^2 = 0$. If $\mathcal{E}$ is an A -module we define on $\Omega(A)$ -module $\mathcal{E}'$ as a tensor product: $\mathcal{E}' = \mathcal{E} \otimes_A \Omega(A)$ where $\Omega(A)$ is considered as $(A, \Omega(A))$ -bimodule. We can define a connection on A -module $\mathcal{E}$ as

a connection of $\Omega(A)$ -module $\mathcal{E}'$; this definition is equivalent to the definition given by Connes (see[1]).

In this definition of connection on A -module $\mathcal{E}$ the algebra $\Omega(A)$ can be replaced with any differential extension of the algebra A (with any differential graded algebra Ω that contains A as a subalgebra of Ω^0). Moreover, one can consider any Q -extension of A (any Q -algebra Ω obeying $A \subset \Omega^0$) and define a connection on A -module E as a connection on Ω -module $E \otimes_A \Omega$. It is interesting to notice that under certain conditions on algebra Ω any projective Ω -module $\mathcal{E}$ can be represented in the form $E \otimes_A \Omega$ where E is projective A -module, $A = \Omega^0$ (see[11]). In particular, this statement is correct if $\Omega = \sum_{0 \leq k \leq n} \Omega^k$ (i.e. the degree of an element of Ω is non-negative and bounded from above).

If a Lie algebra L acts on A by means of infinitesimal automorphisms (derivations) we can construct a differential graded algebra $\Omega = \Omega(L, A)$ of cochains of Lie algebra L with values in A . The elements of Ω can be considered as A -valued functions of anticommuting variables $c^1, \dots, c^n$ corresponding to the elements of the basis $\delta_1, \dots, \delta_n \in L$; the differential d has the form

$$d\omega = (\delta_\alpha \omega) c^\alpha + \frac{1}{2} f_{\beta\gamma}^\alpha c^\beta c^\gamma \frac{\partial}{\partial c^\alpha}$$

where $f_{\beta\gamma}^\alpha$ are the structure constants of L in the basis $\delta_1, \dots, \delta_n$.

In other words we can describe the vector space $\Omega(L, A)$ as a tensor product $\Lambda(L^*) \otimes A$ where L^* stands the vector space dual to L and $\Lambda(M)$ denotes the Grassmann algebra generated by vector space M (as vector space $\Lambda(M)$ is a direct sum of antisymmetric tensor powers of M). The grading on $\Omega(L, A)$ is defined by means of the natural grading on $\Lambda(L^*)$; if A is a graded algebra one should take into account also the grading on A .

Let us consider in more detail connections on A -module E with respect to differential extension $\Omega = \Omega(L, A)$. In this case $\mathcal{E}^0 = E \otimes_A \Omega^0 = E$, $\mathcal{E}^1 = E \otimes_A \Omega^1 = E \otimes_{\mathbb{C}} L^*$. The elements $e \otimes \omega$, $e \in E$, $\omega \in \Omega$ span $\mathcal{E}$, therefore, the connection $\nabla : \mathcal{E}^r \rightarrow \mathcal{E}^{r+1}$ is completely determined by the map $\nabla : \mathcal{E}^0 \rightarrow \mathcal{E}^1$ that can be considered as a map $\nabla : E \rightarrow E \otimes L^*$ or as a family of maps $\nabla_x : E \rightarrow E$ that depend linearly on $x \in L$. Instead of the family ∇_x we can consider n maps $\nabla_1, \dots, \nabla_n$ corresponding to the elements of the basis $f_1, \dots, f_n$ of the

Lie algebra L . These maps obey the Leibniz rule

$$\nabla_\alpha(ea) = \nabla_\alpha e \cdot a + e\delta_\alpha a$$

where δ_α stands for the derivation of the algebra A that corresponds to $f_\alpha \in L$.

Let A be an algebra A_θ of smooth functions on d -dimensional noncommutative torus (i.e. an algebra of expressions of the form $\sum c_n U_n$, where c_n is a $\mathbf{C}$ -valued function on a d -dimensional lattice that vanishes at infinity faster than any power and the multiplication is defined by the formula $U_n U_m = \exp(\pi i \theta_{nm}) U_{n+m}$, where θ_{nm} is a bilinear function on the lattice). Then it is natural to construct a differential extension of A_θ taking as L the Lie algebra of derivations δ_x where $\delta_x U_l = \langle x, l \rangle U_l$. (We assume that the lattice is embedded into vector space V . The vector x belongs to the dual space V^* that can be identified with the Lie algebra L .) Connections corresponding to this differential extension of A_θ appear naturally in the study of toroidal compactifications of M(atric) theory.

Let us suppose now that in addition to the action of Lie algebra on A we have a finite group G acting on A and L by means of automorphisms and that actions of G on A and L are compatible. (If we denote automorphisms of A and of L corresponding to the element $\gamma \in G$ by the same letter γ this means that $\gamma(T(a)) = (\gamma T) \cdot (\gamma a)$ for every $\gamma \in G$, $T \in L$, $a \in A$.) One can define in natural way the action of G on the algebra $\Omega = \Omega(L, A)$; this action commutes with the differential. This means that we can regard the crossed product $\Omega \rtimes G$ as a differential algebra; we have $(\Omega \rtimes G)_0 = \Omega_0 \rtimes G = A \rtimes G$ and therefore the crossed product can be considered as differential extension of $A \rtimes G$.

An $A \rtimes G$ -module E can be considered as an A -module equipped with action of the group G that is compatible with the action of G on A (more precisely we should have $\gamma(xa) = \gamma(x) \cdot \gamma(a)$). As always a connection on E is defined as a connection ∇ on $\Omega \rtimes G$ -module

$$\mathcal{E} = E \otimes_{A \rtimes G} (\Omega \rtimes G).$$

Again this connection is completely determined by the map $\nabla : \mathcal{E}^0 \rightarrow \mathcal{E}^1$ that can be considered as a map

$$\nabla : E \rightarrow E \otimes_{A \rtimes G} (\Omega^1(L, A) \rtimes G)$$

or as a map

$$\nabla : E \rightarrow E \otimes L^*$$

that determines a connection on A -module E and is compatible with the action of the group G on E and on $E \otimes L^*$.

In the case when A is an algebra of functions on noncommutative torus the connections we obtained are precisely the connections that arise by compactification of M(atric) theory on toroidal orbifolds (see [9],[10]).

Conclusion.

In present paper we generalized the theory of connections on modules over associative algebra. We embedded this theory into the theory of connections on modules over Q -algebras and proved a general duality theorem in this framework. Namely, we proved that under certain conditions there exists one-to-one correspondence between connections on modules over one Q -algebra and connections on modules over another Q -algebra and found relation between corresponding curvatures. More precisely, it follows from our results that under certain conditions gauge theory constructed by means of Q -algebra Ω is equivalent to gauge theory corresponding to the Q -algebra $\text{End}_\Omega \mathcal{E}$ where $\mathcal{E}$ is an Ω -module equipped with a connection. (We use the fact that every connection determines a structure of Q -algebra on the algebra of endomorphisms.) This theorem can be applied to many concrete situations; we are planning to give applications in forthcoming papers.

Acknowledgments.

I am very grateful to M. Khovanov, M. Kontsevich, M. Rieffel and D. Sternheimer for useful comments. I am deeply indebted to Caltech Theory Group, especially to J. Schwarz and E. Witten, for warm hospitality and interesting discussions.

Appendix

Differential algebras, A_∞ -algebras and Q -algebras.

If Ω is a Q -algebra one can extend it to a differential algebra Ω' adjoining new element X obeying $X^2 = \omega$, $a_1 X a_2 = 0$ for all $a_1, a_2 \in \Omega$. The differential on Ω' is defined by the formula $d\omega = Q\omega + [X, \omega]$ (this construction was used by A. Connes [1]). The differential algebra we constructed is equivalent in some sense to the Q -algebra we started with.

One can define an A_∞ -algebra as a vector space V equipped with multilinear operations m_i ; these operations should satisfy some relations. (The operations m_i determine a derivation of tensor algebra over V ; the square of this derivation should be equal to

zero.) In standard definition of A_∞ -algebra one considers operations m_i where the number of arguments i is ≥ 1 . However, one can modify the definition including an operation m_0 (if the number of arguments is equal to zero, then the operation is simply a fixed element of V). Using the modified definition one can say that Q -algebra is an A_∞ -algebra where all operations with the number of arguments ≥ 3 vanish. (In standard definition this requirement leads to differential algebras.)

References.

1. Connes, A., *Noncommutative Geometry*, Academic Press, New York, (1994).
2. Karoubi, M., *Homology cyclique et K-theorie*, Asterisque, 149 (1987), 147 pp.
3. Kastler, D., *Cyclic cohomology within the differential envelope*, Hermann, Paris, (1988), 184 pp.
4. Connes, A., Douglas, M., and Schwarz, A., *Noncommutative geometry and Matrix theory: compactification on tori*, JHEP **02** (1998) 003; hep-th/9711162.
5. Schwarz, A., *Morita equivalence and duality*, Nucl. Phys. **B534** (1998) pp. 720-738; hep-th/9805034.
6. Konechny, A. and Schwarz, A., *BPS states on noncommutative tori and duality*, Nucl. Phys. **B550** (1999) 561-584; hep-th/9811159.
- Konechny, A. and Schwarz, A., *Supersymmetry algebra and BPS states of super Yang-Mills theories on noncommutative tori*, Phys. Lett. **B453** (1999) 23-29; hep-th/9901077.
7. Konechny, A. and Schwarz, A., *$1/4$ BPS states on noncommutative tori*, JHEP **09** (1999) 030; hep-th/9907008.
8. Nekrasov, N. and Schwarz, A., *Instantons on noncommutative R^4 , and $(2,0)$ superconformal six-dimensional theory*, hep-th/9802068.
9. Seiberg, N. and Witten, E., *String Theory and Noncommutative Geometry*, JHEP **9909** (1999) 032; hep-th/9908142.
10. Ho, P.-M. and Wu, Y.-S., *Noncommutative Gauge Theories in Matrix Theory*, Phys.Rev. **D58** (1998) 066003; hep-th/9801147.
11. Konechny, A. and Schwarz, A., *Compactification of M (atrix) theory on noncommutative toroidal orbifolds*, hep-th/9912185
12. Bass, H., *Algebraic K-theory*, Benjamin, NY-Amsterdam, (1962), 762 pp

12. Schwarz, A. *Geometry of Batalin-Vilkovisky quantization*, CMP, **155** (1993) 249-260, *Semiclassical approximation in Batalin-Vilkovisky formalism*, CMP,**158** (1993) 373-396, Alexandrov,M., Kontsevich, M., Schwarz, A. and Zaboronsky, O. ,*The geometry of master equation and topological quantum field theory*, Int. J. of Modern Physics,**A12** (1997) 1405-1429